

5. Jimbow M, Jimbow K, Kanoh H: Characterization of biochemical properties of melanosomes for structural and functional differentiation: analysis of the compositions of lipids and proteins in melanosomes and their subfractions.
6. Jimbow K, Quevedo WC Jr, Fitzpatrick TB, Szabo G: Some aspects of melanin biology: 1950-1975. *J Invest Dermatol* 67:72-89, 1976

7. Jimbo K, Fitzpatrick TB, Quevedo WC Jr: Cascade of melanogenesis in epidermal melanin pigmentation: the melanosome as a programmed organelle. *Struct Funct Melanin* 2:71-82, 1985

HNK-1 Antigen Is not Specific for Natural Killer Cells

To the Editor:

we read with great interest the article by Habets et al, published in the March 1988 issue [1]. The peritumoral inflammatory infiltrate of basal cell carcinoma (BCC) has been investigated by using a series of monoclonal antibodies. The authors provided evidence for a minor participation of natural killer (NK) cells, in defense against BCC, on the basis of the low percentage of Leu-7 (HNK-1)-positive cells in the infiltrate. Unfortunately the HNK-1 antigen, isolated by Abo and Balch in 1981 [2], cannot be considered specific for NK cells. Rather, the HNK-1 antigen is coexpressed by most CD8-positive/CD11b-positive suppressor cells [3]. Indeed, Phillips and Babcock identified a new antigen in 1983 [4], named NKP-15 (CD16), considered to be specific for essentially all human NK cells. A number of findings demonstrate that CD16 antigen is specific for NK cells, while the HNK-1 antigen is not. First, functional studies from Lanier et al [5] and Abo et al [6] have well established that CD16-positive cells display high levels of NK activity, while HNK-1-positive/CD16-negative cells possess low levels of cytotoxic cell function. On the other hand, CD16-positive/HNK-1-negative cells, when stimulated with interleukin-2 [7] or with NK-sensitive tumor cells K-562 [8], markedly augment their NK cell activity, while stimulated HNK-1-positive/CD16-negative cells never display higher NK cell function. Moreover, CD16-positive cells are able to phagocytize AET-sheep red blood cells (SRBC), whereas HNK-1-positive/CD16-negative cells never display phagocytic capability for AET-SRBC [9]. As for morphologic characteristics, CD16-positive cells show significant ultrastructural differences in comparison to HNK-1-positive/CD16-negative cells [10,11], as previously demonstrated by using a peroxidase-colloidal gold double labeling in immunoelectron microscopy [12]. Ultrastructural differences were confirmed by morphometric investigations [13]. Finally, a new antigen, named NKH-1 (Leu-19), which is expressed on the whole non-major histocompatibility complex-restricted cytotoxic cell population, comprising NK cells has recently been described [14,15].

In conclusion, the HNK-1-positive/CD16-negative cell subset seems to be distinct from the CD16-positive NK cell population. As a result, immunohistochemical studies performed by Werner Habets et al cannot rule out the possibility that NK cells play a role in the defense against BCC. This issue could be resolved by further investigations dealing with CD16 and NKH-1 antigens.

Gian Carlo Manara, M.D.
Corrado Ferrari, M.D.
Giuseppe De Panfilis, M.D.
University of Parma
Parma, Italy

REFERENCES

1. Habets JMW, Tank B, Vuzevski VD, van Reede EC, Stolz E, van Joost T: Characterization of the mononuclear infiltrate in basal cell carcinoma: a predominantly T cell-mediated immune response with minor participation of Leu-7⁺ (natural killer) and Leu-14⁺ (B) cells. *J Invest Dermatol* 90:289-292, 1988
2. Abo T, Balch CM: A differentiation antigen of human NK and K cells identified by a monoclonal antibody (HNK-1). *J Immunol* 127:1024-1029, 1981

3. Clement LT, Grossi CE, Gartland GL: Morphologic and phenotypic features of the subpopulation of Leu-2⁺ cells that suppresses B cell differentiation. *J Immunol* 133:2461-2468, 1984
4. Phillips JH, Babcock GF: NKP-15: a monoclonal antibody reactive against purified natural killer cells and granulocytes. *Immunol Lett* 6:143-149, 1983
5. Lanier LL, Le AM, Phillips JH, Warner NL, Babcock GF: Subpopulations of human natural killer cells defined by expression of the Leu-7 (HNK-1) and Leu-11 (NK-15) antigens. *J Immunol* 131:1789-1796, 1983
6. Abo T, Miller CA, Balch CM: Characterization of human granular lymphocyte subpopulations expressing HNK-1 (Leu-7) and Leu-11 antigens in the blood and lymphoid tissues from fetuses, neonates and adults. *Eur J Immunol* 14:616-623, 1984
7. Lanier LL, Benike CJ, Phillips JH, Engleman EG: Recombinant interleukin 2 enhanced natural killer cell-mediated cytotoxicity in human lymphocyte subpopulations expressing the Leu-7 and Leu-11 antigens. *J Immunol* 134:794-801, 1985
8. Phillips JH, Lanier LL: A model for the differentiation of human natural killer cells. Studies on the in vitro activation of Leu-11⁺ granular lymphocytes with a natural killer-sensitive tumor cell, K562. *J Exp Med* 161:1464-1482, 1985
9. Manara GC, Sansoni P, Ferrari C, De Panfilis G: Natural killer cells expressing the Leu-11 antigen display phagocytic activity for 2-aminoethylisothiourea bromide hydrobromide-treated sheep red blood cells. *Lab Invest* 55:412-418, 1986
10. Manara GC, De Panfilis G, Ferrari C: Ultrastructural characterization of human large granular lymphocyte subsets defined by the expression of HNK-1 (Leu-7), Leu-11, or both HNK-1 and Leu-11 antigens. *J Histochem Cytochem* 33:1129-1133, 1985
11. Manara GC, Arancia G, Fiorentini C, Ferrari C, De Panfilis G: Ultrastructural differences between Leu-7⁺, 11⁻ NK cells and Leu-11⁺ NK cells. *J Immunol Meth* 98:155-156, 1987
12. Manara GC, De Panfilis G, Ferrari C, Scandroglio R: Immunoperoxidase-immunogold double labelling in immunoelectron microscopy of large granular lymphocytes. *J Immunol Meth* 75:189-192, 1983
13. Arancia G, Fiorentini C, Ferrari C, De Panfilis G, Manara GC: Morphometric characterization of NK cell subset expressing the Leu-11 antigen in comparison to Leu-7 positive, 11 negative cells. *Cell Biol Int Rep* 10:845-853, 1986
14. Lanier LL, Le AM, Civin CI, Loken MR, Phillips JH: The relationship of CD16 (Leu-11) and Leu-19 (NKH-1) antigen expression on human peripheral blood NK cells and cytotoxic T lymphocytes. *J Immunol* 136:4480-4486, 1986
15. Manara GC, Ferrari C, Sansoni P, De Panfilis G: Immunoelectron microscopic characterization of human natural killer cells related to the expression of CD16 and Leu-19 antigens (abstr). *J Invest Dermatol* 89:331, 1987

REPLY

The authors acknowledge Dr. Manara's comments regarding our recent publication [1]. Indeed, as stated in our article we did not totally exclude the participation of leu 7 (HNK-1)-positive cells in the defense against Basal cell carcinoma (BCC). In a recent publication Kohchiyama et al [2], using leu 7 (HNK-1) monoclonal antibody, also concluded that the cellular cytotoxicity mediated by NK cells may not be the main defense against BCC. Although HNK-1 antigen may be coexpressed by most leu 2a (CD-8) positive suppres-